

Grade A and B Recommendations of the United States Preventive Services Task Force

Topic	Text	Grade	Date In Effect
Screening for abdominal aortic aneurysm	The USPSTF recommends one-time screening for abdominal aortic aneurysm (AAA) by ultrasonography in men aged 65 to 75 who have ever smoked.	B	Feb 28, 2005
Screening and counseling to reduce alcohol misuse	The U.S. Preventive Services Task Force (USPSTF) recommends screening and behavioral counseling interventions to reduce alcohol misuse (go to Clinical Considerations) by adults, including pregnant women, in primary care settings.	B	April 30, 2004
Aspirin to prevent CVD: men	The USPSTF recommends the use of aspirin for men age 45 to 79 years when the potential benefit due to a reduction in myocardial infarctions outweighs the potential harm due to an increase in gastrointestinal hemorrhage.	A	March 30, 2009
Aspirin to prevent CVD: women	The USPSTF recommends the use of aspirin for women age 55 to 79 years when the potential benefit of a reduction in ischemic strokes outweighs the potential harm of an increase in gastrointestinal hemorrhage.	A	March 30, 2009
Screening for bacteriuria	The USPSTF recommends screening for asymptomatic bacteriuria with urine culture for pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	A	July 31, 2008
Screening for high blood pressure	The U.S. Preventive Services Task Force (USPSTF) recommends screening for high blood pressure in adults aged 18 and older.	A	Dec 31, 2007
Counseling related to	The USPSTF recommends that women whose	B	Sept 30, 2005

BRCA screening	family history is associated with an increased risk for deleterious mutations in BRCA1 or BRCA2 genes be referred for genetic counseling and evaluation for BRCA testing.		
Screening for breast cancer (mammography)	The USPSTF recommends screening mammography for women with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older.	B	September 30, 2002
Chemoprevention of breast cancer	The USPSTF recommends that clinicians discuss chemoprevention with women at high risk for breast cancer and at low risk for adverse effects of chemoprevention. Clinicians should inform patients of the potential benefits and harms of chemoprevention.	B	July 31, 2002
Interventions to support breast feeding	The USPSTF recommends interventions during pregnancy and after birth to promote and support breastfeeding.	B	Oct 31, 2008
Screening for cervical cancer	The USPSTF strongly recommends screening for cervical cancer in women who have been sexually active and have a cervix.	A	Jan 31, 2003
Screening for chlamydial infection: non-pregnant women	The U.S. Preventive Services Task Force (USPSTF) recommends screening for chlamydial infection for all sexually active non-pregnant young women aged 24 and younger and for older non-pregnant women who are at increased risk.	A	June 30, 2007
Screening for chlamydial infection: pregnant women	The USPSTF recommends screening for chlamydial infection for all pregnant women aged 24 and younger and for older pregnant women who are at increased risk.	B	June 30, 2007
Screening for	The U.S. Preventive Services Task Force (USPSTF)	A	June 30,

cholesterol abnormalities: men 35 and older	strongly recommends screening men aged 35 and older for lipid disorders.		2008
Screening for cholesterol abnormalities: men younger 35	The USPSTF recommends screening men aged 20 to 35 for lipid disorders if they are at increased risk for coronary heart disease.	B	June 30, 2008
Screening for cholesterol abnormalities: women 45 and older	The USPSTF strongly recommends screening women aged 45 and older for lipid disorders if they are at increased risk for coronary heart disease.	A	June 30, 2008
Screening for cholesterol abnormalities: women younger than 45	The USPSTF recommends screening women aged 20 to 45 for lipid disorders if they are at increased risk for coronary heart disease.	B	June 30, 2008
Screening for colorectal cancer	The USPSTF recommends screening for colorectal cancer (CRC) using fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults, beginning at age 50 years and continuing until age 75 years. The risks and benefits of these screening methods vary.	A	Oct 31, 2008
Chemoprevention of dental caries	The USPSTF recommends that primary care clinicians prescribe oral fluoride supplementation at currently recommended doses to preschool children older than 6 months of age whose primary water source is deficient in fluoride.	B	April 30, 2004
Screening for depression: adults	The USPSTF recommends screening adults for depression when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up.	B	Dec 31, 2009

Screening for depression: adolescents	The USPSTF recommends screening of adolescents (12-18 years of age) for major depressive disorder (MDD) when systems are in place to ensure accurate diagnosis, psychotherapy (cognitive-behavioral or interpersonal), and follow-up.	B	March 30, 2009
Screening for diabetes	The USPSTF recommends screening for type 2 diabetes in asymptomatic adults with sustained blood pressure (either treated or untreated) greater than 135/80 mm Hg.	B	June 30, 2008
Counseling for a healthy diet	The USPSTF recommends intensive behavioral dietary counseling for adult patients with hyperlipidemia and other known risk factors for cardiovascular and diet-related chronic disease. Intensive counseling can be delivered by primary care clinicians or by referral to other specialists, such as nutritionists or dietitians.	B	Jan 30, 2003
Supplementation with folic acid	The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 g) of folic acid.	A	May 31, 2009
Screening for gonorrhea: wp,em	The U.S. Preventive Services Task Force (USPSTF) recommends that clinicians screen all sexually active women, including those who are pregnant, for gonorrhea infection if they are at increased risk for infection (that is, if they are young or have other individual or population risk factors; go to Clinical Considerations for further discussion of risk factors).	B	May 31, 2005
Prophylactic medication for	The USPSTF strongly recommends prophylactic ocular topical medication for all newborns against	A	May 31, 2005

gonorrhoea: newborns	gonococcal ophthalmia neonatorum.		
Screening for hearing loss	The USPSTF recommends screening for hearing loss in all newborn infants.	B	July 31, 2008
Screening for hemoglobinopathies	The U. S. Preventive Services Task Force (USPSTF) recommends screening for sickle cell disease in newborns.	A	Sept 30, 2007
Screening for hepatitis B	The U.S. Preventive Services Task Force (USPSTF) strongly recommends screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit.	A	June 30, 2009
Screening for HIV	The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians screen for human immunodeficiency virus (HIV) all adolescents and adults at increased risk for HIV infection (go to Clinical Considerations for discussion of risk factors).	A	July 31, 2005
Screening for congenital hypothyroidism	The USPSTF recommends screening for congenital hypothyroidism (CH) in newborns.	A	March 31, 2008
Screening for iron deficiency anemia	The USPSTF recommends routine screening for iron deficiency anemia in asymptomatic pregnant women.	B	May 31, 2006
Iron supplementation in children	The U.S. Preventive Services Task Force (USPSTF) recommends routine iron supplementation for asymptomatic children aged 6 to 12 months who are at increased risk for iron deficiency anemia (go to Clinical Considerations for a discussion of increased risk).	B	May 30, 2006
Screening and	The USPSTF recommends that clinicians screen	B	Dec 31, 2003

counseling for obesity: adults	all adult patients for obesity and offer intensive counseling and behavioral interventions to promote sustained weight loss for obese adults.		
Screening and counseling for obesity: children	The USPSTF recommends that clinicians screen children aged 6 years and older for obesity and offer them or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.	B	Jan 31, 2010
Screening for osteoporosis	The U.S. Preventive Services Task Force (USPSTF) recommends that women aged 65 and older be screened routinely for osteoporosis. The USPSTF recommends that routine screening begin at age 60 for women at increased risk for osteoporotic fractures. (Go to Clinical Considerations for discussion of women at increased risk.)	B	Sept 30, 2002
Screening for PKU	The USPSTF recommends screening for phenylketonuria (PKU) in newborns.	A	March 31, 2008
Screening for Rh incompatibility: first pregnancy visit	The U.S. Preventive Services Task Force (USPSTF) strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.	A	Feb 29, 2004
Screening for Rh incompatibility: 24-28 weeks gestation	The USPSTF recommends repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	B	Feb 29, 2004
Counseling for STIs	The USPSTF recommends high-intensity behavioral counseling to prevent sexually transmitted infections (STIs) for all sexually active adolescents and for adults at increased risk for	B	Oct 31, 2008

STIs.

Screening for syphilis: non-pregnant persons	The U.S. Preventive Services Task Force (USPSTF) strongly recommends that clinicians screen persons at increased risk for syphilis infection.	A	July 31, 2004
Screening for syphilis: pregnant women	The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.	A	July 31, 2004
Counseling for tobacco use	The USPSTF recommends that clinicians ask all adults about tobacco use and provide tobacco cessation interventions for those who use tobacco products.	A	April 30, 2009
Counseling for tobacco use	The USPSTF recommends that clinicians ask all pregnant women about tobacco use and provide augmented, pregnancy-tailored counseling for those who smoke.	A	April 30, 2009
Screening for visual acuity in children	The USPSTF recommends screening to detect amblyopia, strabismus, and defects in visual acuity in children younger than age 5 years.		